

Review

Ying Dong, Brittney S. Harrington, Mark N. Adams, Andreas Wortmann, Sally-Anne Stephenson, Jessica Lisle, Adrian Herington, John D. Hooper and Judith A. Clements*

Activation of membrane-bound proteins and receptor systems: a link between tissue kallikrein and the KLK-related peptidases

Abstract: The 15 members of the kallikrein-related serine peptidase (KLK) family have diverse tissue-specific expression profiles and roles in a range of cellular processes, including proliferation, migration, invasion, differentiation, inflammation and angiogenesis that are required in both normal physiology as well as pathological conditions. These roles require cleavage of a range of substrates, including extracellular matrix proteins, growth factors, cytokines as well as other proteinases. In addition, it has been clear since the earliest days of KLK research that cleavage of cell surface substrates is also essential in a range of KLK-mediated cellular processes where these peptidases are essentially acting as agonists and antagonists. In this review we focus on these KLK-regulated cell surface receptor systems including bradykinin receptors, proteinase-activated receptors, as well as the plasminogen activator, ephrins and their receptors, and hepatocyte growth factor/Met receptor systems and other plasma membrane proteins. From this analysis it is clear that in many physiological and pathological settings KLKs have the potential to regulate multiple receptor systems simultaneously; an important issue when these peptidases and substrates are targeted in disease.

Keywords: cell membrane-bound protein; kallikrein-related peptidase; receptor; signalling pathways, tissue kallikrein.

DOI 10.1515/hsz-2014-0147

Received February 17, 2014; accepted April 12, 2014; previously published online April 18, 2014

Introduction

Tissue kallikrein or kallikrein 1 (*KLK1/KLK1*) was the first described member of a gene family located at human chromosome 19q13.4 that encodes 15 serine proteinases named kallikrein-related peptidases (KLKs) (Gan et al., 2000; Harvey et al., 2000; Yousef and Diamandis, 2001). The action of KLK1, the archetypal member of this family, was first recognised over 100 years ago in the work of Abelous and Bardier who discovered that an alcohol-insoluble fraction of human urine caused peripheral vasodilation and hypotension in dogs (Abelous and Bardier, 1909). Later studies showed that the hypotensive effect occurs, at least in part, by the action on bradykinin receptors, a sub-family of G protein-coupled receptors, of kinin peptides released from kininogens by KLK1 (Werle, 1934; Rocha e Silva et al., 1949; Bhoola et al., 1992). Notably, it became clear from the earliest days of research on the extended KLK family that these KLK1-related secreted extracellular peptidases also can have important physiological and pathological effects via actions on cell surface proteins. As summarised in Figure 1 and Table 1, it is now clear that these effects can occur both indirectly, for example by processing of precursor ligands as is the case of KLK1 cleavage of kininogens; and directly, for example by proteolysis of cell surface proteins as has been reported also to occur when bradykinin receptors are cleaved by KLK1. This theme has certainly been borne out for other members of the KLK family with many examples reported over the last 20 years

*Corresponding author: Judith A. Clements, Cancer Program, Institute of Health and Biomedical Innovation, Queensland University of Technology, Translational Research Institute, 37 Kent Street, Woolloongabba 4102, Queensland, Australia, e-mail: j.clements@qut.edu.au

Ying Dong, Sally-Anne Stephenson, Jessica Lisle and Adrian

Herington: Cancer Program, Institute of Health and Biomedical Innovation, Queensland University of Technology, Translational Research Institute, 37 Kent Street, Woolloongabba 4102, Queensland, Australia

Brittney S. Harrington, Mark N. Adams, Andreas Wortmann and John D. Hooper: Mater Medical Research Institute, The University of Queensland, Translational Research Institute, 37 Kent Street, Woolloongabba 4102, Queensland, Australia




Figure 1 Schematic diagram showing KLKs 1, 2 and 4-8 activation of receptors and/or membrane-bound factors initiating their downstream signalling pathways.

BK, bradykinin; BKR, bradykinin receptors 1 and 2; CDP1, CUB domain containing protein; HGF(A), hepatocyte growth factor (activator); KG, kininogen; KLK1, tissue kallikrein; PAR, proteinase-activated receptor; uPA(R), urokinase plasminogen activator (receptor).

of KLK-mediated phenotypes that require activation of cell surface receptor systems. In the following sections we summarise these findings with a particular focus on bradykinin receptors, proteinase-activated receptors, as well as the plasminogen activator and hepatocyte growth factor/Met receptor systems and other cell surface proteins.

Kallikrein-kinin system

More than 100 years ago, Abelous and Bardier began to elucidate an important signalling pathway that has come to be known as the kallikrein-kinin system. The activating enzymes for this system are plasma kallikrein and tissue kallikrein, which are derived from two unrelated genes (Clements, 2000). The focus of this review is tissue kallikrein or KLK1. Tissue kallikrein (KLK1), the founding member of the kallikrein-related peptidase (KLK) family (Kraut et al., 1930), is a glycoprotein with trypsin-like substrate specificity for cleavage after arginine (Arg) and lysine (Lys) residues. It is expressed in a range of tissues and organs, and can process kininogens, growth factors, and extracellular matrix (ECM) molecules to regulate inflammation and pain, blood vessel dilation and permeability, blood pressure and sodium homeostasis control, kidney, heart and neurological function (Fiedler, 1979; Schachter, 1979; Brady et al., 1989; Clements, 2000; Chao et al., 2006; Madeddu et al., 2007; Savvatis et al., 2010; Rhaleb et al., 2011; Stadnicki, 2011; Viel and Buck, 2011; Kayashima et al., 2012; Regoli et al., 2012; da Costa et al., 2014). As an extracellular peptidase, KLK1 exerts

these pleotropic biological effects by cleavage of the substrate, low molecular weight (LMW) kininogen, generating the kinin, lys-bradykinin, which exerts its biological functions via binding to the G protein-coupled receptor, the bradykinin B2 (Marcondes and Antunes, 2005). The related bradykinin B1 receptor is predominantly activated by a cleaved degradation product of lys-bradykinin at its C-terminal Arg by kininase I, lys-des-arg-bradykinin. The B1 receptor is largely an inducible receptor up-regulated in inflammatory conditions and tissue injury; conversely, the B2 receptor is constitutively expressed in a diverse range of healthy tissues mediating the action of kinins as well as up-regulated in disease (Madeddu et al., 2007; Savvatis et al., 2010; Stadnicki, 2011; Viel and Buck, 2011; Kayashima et al., 2012; Regoli et al., 2012). KLK1 also directly induces rat uterine contraction and cardio-protection independent of kinin formation (Chao et al., 1981, 2008), further suggesting KLK1 has potential biological functions by acting on other substrates in addition to kininogen. A recent study showed that migration of circulating proangiogenic cells induced by KLK1 may not be all mediated by kinin *in vitro* while kinins are involved in the function of KLK1 *in vivo* (Spinetti et al., 2011). Of particular interest to this review, the human bradykinin B2 receptor can be activated in kininogen-free systems in which cells are exposed to KLK1 and trypsin, but the direct cleavage of the B2 receptor to generate a signal has yet to be observed (Hecquet et al., 2000; Biyashev et al., 2006). It thus remains to be seen if the B2 receptor can be considered as a ‘proteolytically-activated receptor’ akin to the proteinase-activated receptors (PARs), which are

Table 1 Summary of activation of cell membrane proteins and receptors by KLKs.

Kallikreins	Membrane proteins/receptors and signalling systems	References
KLK1	Kininogen: KLK1-(brady)kinin system (Brady)kinin receptor (BKR)2 PAR1: PAR1-PKC-Src-MMP, keratinocyte migration; migration of prostate cancer cells BKR2, migration of circulating proangiogenic cells	(Marcondes and Antunes, 2005) (Biyashev et al., 2006) (Gao et al., 2010a) (Gao et al., 2010b) (Spinetti et al., 2011)
KLK2	uPA	(Frenette et al., 1997)
KLK4	PAR1, PAR2, uPA-uPAR pro-meparin β EphB4 HGF/MET CDCP1	(Mize et al., 2008; Ramsay et al., 2008; Gratio et al., 2010) (Takayama et al., 2001; Beaufort et al., 2006) (Becker-Pauly et al., 2007; Ohler et al., 2010) (This review) (Mukai et al., 2008) This review
KLK5	Pro-HGFA PAR2 Desmoglein 1, cell-cell junction in oral squamous cancer	(Mukai et al., 2008) (Stefansson et al., 2008) (Jiang et al., 2011)
KLK6	Amyloid precursor protein (APP) in Alzheimer's disease and Parkinson's disease; PAR1, PAR2, PAR4	(Little et al., 1997; Ogawa et al., 2000; Magklara et al., 2003) (Oikonomopoulou et al., 2006)
KLK7	Desmocollin, corneodesmosin; E-cadherin uPAR	(Caubet et al., 2004) (Johnson et al., 2007) (Ramani and Haun, 2008a)
KLK8	rPAR2 but not hPAR2	(Stefansson et al., 2008; Ramachandran et al., 2012)
KLK14	PAR2	(Stefansson et al., 2008; Gratio et al., 2011; Chung et al., 2012)

activated by a proteolytically-unmasked tethered ligand (Adams et al., 2011). These data have for the first time shown that a KLK peptidase could activate a membrane-bound receptor and led the way for many of the studies described below.

Proteinase-activated receptors

PARs are G protein-coupled receptors – PAR1 to PAR4 – that are activated by proteolysis primarily mediated by serine proteinases with trypsin-like specificity for cleavage after Arg and Lys, including members of the KLK family (Vu et al., 1991; Bohm et al., 1996; Ishihara et al., 1997; Xu et al., 1998; Boire et al., 2005; Vesey et al., 2007) as shown diagrammatically for KLK1 and KLK4 in Figure 1. These proteolytic agonists remove the PAR extracellular amino terminal pro-domain, unmasking a neo-epitope, termed the tethered ligand, that binds intramolecularly to induce intracellular signal transduction. Via these mechanisms, PARs function as plasma membrane sensors of extracellular and cell surface associated proteinases that regulate cellular responses such as proliferation, migration, inflammatory responses, thrombus formation and angiogenesis, and contribute extensively to homeostasis

as well as to disease progression (Macfarlane et al., 2001; Adams et al., 2011).

Proteolytic activation of human PAR1, occurring on the carboxyl side of residue Arg⁴¹, can be mediated by KLK1 (Gao et al., 2010a,b), KLK4 (Mize et al., 2008; Ramsay et al., 2008; Gratio et al., 2010), and KLK5, 6 and 14 (Oikonomopoulou et al., 2006). Similarly, PAR2 activation after Arg³⁶ can occur through proteolysis by KLK2 (Mize et al., 2008), KLK4 (Mize et al., 2008; Ramsay et al., 2008), KLK5, KLK6 (Oikonomopoulou et al., 2006) and KLK14 (Oikonomopoulou et al., 2006; Gratio et al., 2010), while PAR4 activation by cleavage following Arg⁴⁷ can be carried out by KLK14 (Oikonomopoulou et al., 2006). Consistently, another study demonstrated that while KLK6 is able to efficiently cleave a peptide spanning the PAR2 activation site, this proteinase could not cleave peptides spanning PAR1, PAR3 or PAR4 activation sites (Angelo et al., 2006). Interestingly, apart from thrombin no other PAR3 targeting proteinases have been identified (Ishihara et al., 1997). It is also important to note that KLKs can function as PAR antagonists by cleavage at sites other than the canonical activation site, leading to receptor inactivation or so-called 'disarming' (Adams et al., 2011). For example, KLK1 can disarm PAR1 (Gao et al., 2010a,b) and KLK14 can inactivate both PAR1 and PAR4 (Oikonomopoulou et al., 2006).

In terms of the functional consequences of KLK activation of PARs, several studies have demonstrated that these mechanisms regulate important cellular responses in both physiological and pathological settings. For example, KLK1 promotes keratinocyte proliferation and migration *in vitro* and in an *in vivo* rat skin wound healing model via a PAR1 dependent, BKR2 independent mechanism requiring downstream activation of PKC and Src resulting in increased matrix metalloproteinase activity (MMP) and regulation of EGFR signalling (Gao et al., 2010a). This group has also recently shown that KLK1, via PAR1, promotes migration and invasion of prostate cancer DU145 but not lung cancer A549 cells (Gao et al., 2010b). Signalling via this PAR and PAR2 induced by both KLK2 and KLK4 enhanced proliferation of prostate cancer DU145 cells (Mize et al., 2008). Interestingly, using murine whole splenocyte preparations and the human Jurkat T cell line it has been shown that KLK6 via PAR1 promotes pro-survival and inhibits pro-apoptotic mechanisms, leading to vigorous splenic T cell survival in the presence of the cell death inducing agents, camptothecin, dexamethasone, staurosporine and Fas-ligand (Scarlsbrick et al., 2011). This KLK can also initiate intracellular signalling in neurons via PAR1 and in astrocytes through both PAR1 and PAR2 (Vandell et al., 2008) and has recently been reported to serve as a molecular trigger via PAR1 of processes involved in development of astrogliosis (Scarlsbrick et al., 2012a). In addition, use of function blocking antibodies and pharmacological approaches demonstrated that KLK4 activates PAR1 but not PAR2 in colon cancer cells with PAR1 signalling promoting migration of these cells via ERK1/2 (Gratio et al., 2010). In contrast, KLK14 signals to HT29 colon cancer cells via PAR2 leading to receptor internalization and proliferation (Gratio et al., 2011). It is also noteworthy that KLK4, secreted by prostate cancer epithelia, can activate PAR1 in surrounding stroma to initiate release of the cytokine interleukin-6 that stimulates cancer cell proliferation in a paracrine loop (Wang et al., 2009).

Hepatocyte growth factor/Met receptor system

Hepatocyte growth factor (HGF) and its receptor MET are required for normal development and homeostasis while aberrant activation of this receptor system, via overexpression, mutation or dysfunctional ligand production and/or activation, contributes to malignant transformation and progression of a variety of cancers. In addition, these changes can often be associated with poor patient

outcome and drug resistance (Trusolino et al., 2010). In its precursor state as pro-HGF, the presence of the pro-domain prevents activation of Met signalling. This pro-domain is released by the action of the Arg/Lys-specific serine proteinase HGF activator (HGFA), which itself must be converted from zymogen (pro-HGFA) to active enzyme by other serine proteinases such as thrombin (Mahtouk et al., 2010). In addition to this thrombin/HGFA/HGF Met receptor activating cascade, it appears that members of the KLK family can also initiate signalling via this receptor system through at least two levels as shown diagrammatically for KLK4 in Figure 1. The first level is via activation of pro-HGFA as it has recently been demonstrated that KLK4 and KLK5, but not KLK2 or KLK3, activate this zymogen at its canonical activation site Arg407 (Mukai et al., 2008). Importantly, KLK4 and KLK5 activated HGFA efficiently activated pro-HGF and induced cell scattering and invasion *in vitro* while these actions were strongly inhibited by HGFA inhibitor, a type 1, integral membrane Kunitz-type serine protease inhibitor that inhibits HGFA and other pro-HGF-activating proteases (Mukai et al., 2008). KLK-mediated regulation of this signalling cascade is also possible at the level of pro-HGF activation as it has been shown that KLK4 is able to activate urokinase-type plasminogen activator (Takayama et al., 2001) which is an efficient activator of pro-HGF (Naldini et al., 1992; Mars et al., 1993).

Plasminogen activator system

The plasminogen activator system is involved in cancer progression and metastasis through various processes, such as cell migration, adhesion and cell cycle regulation. Urokinase-type plasminogen activator (uPA) is a serine protease and its receptor uPAR (CD87) is a three domain (D1D2D3) glycoprotein tethered to the cell membrane with a GPI anchor (Huai et al., 2006). Expression of uPA-uPAR has been associated with prognosis in different cancers, although cleaved uPAR has stronger association with clinical outcome of patients than those with intact forms (Reuning et al., 2003; Rasch et al., 2008). For example, expression of uPAR lacking D1 predicted a poor prognosis in the patients with squamous cell lung cancer (Almasi et al., 2005), suggesting that the cleavage of this receptor has a role in cancer progression. KLK2, KLK4, and KLK8 can activate uPA (Frenette et al., 1997; Takayama et al., 2001; Rajapakse et al., 2005; Beaufort et al., 2006). Indeed, the first identified substrate of KLK4 was uPA (Takayama et al., 2001), which can also activate the metalloprotease (MMP) system via plasmin (Schmalfeldt et al., 2001; Smith

and Marshall, 2010), further linking these three proteolytic systems. Interestingly, a later study that examined the proteolytic effects of KLK4 at different concentrations (Beaufort et al., 2006) demonstrated that low levels of KLK4 cleaved uPA while higher KLK4 levels also had an effect on its receptor, uPAR (see Figure 1). This latter proteolytic action generates a soluble D1 fragment and D2D3 fragment still bound to the cell membrane. Beaufort et al. (2006) also showed that KLK4 cleaved uPAR on the membrane of U937 monocytes led to a reduced binding capacity to uPA. It is known that uPAR interacts with integrins and the ECM protein vitronectin (Smith and Marshall, 2010), and the function of the cell membrane-bound uPAR D2D3 domain is in binding to these adhesive proteins, promoting migration of cells and cancer progression. Interestingly, KLK7 is also suggested to play a role in the shedding of uPAR generating a soluble form of this receptor and reduced pancreatic cancer cell adhesion to vitronectin (Ramani and Haun, 2008b). Similar cell biology effects of the proteolytic action of KLK2, KLK4 and KLK8 on uPA-uPAR remain to be studied. Indeed, more in depth studies of the mechanisms involving these two protease systems are required to fully understand their interactive roles in human diseases.

EphB4 and its ligand ephrin-B2

The Eph receptor tyrosine kinases are the largest family of receptor tyrosine kinases with 14 members in two subfamilies designated A and B (Arvanitis and Davy, 2012). Eph receptors are involved in many cellular and developmental processes including development, homeostasis and disease (Bush and Soriano, 2012). Several Eph receptors are also over-expressed in cancer and novel anti-cancer strategies that target these proteins are in development (Nievergall et al., 2012; Boyd et al., 2014). EphB4, in particular, is commonly over-expressed in most epithelial cancers, including 66% of prostate cancers (PCa), 58% of breast cancers and 86% of ovarian cancers (Lee et al., 2005; Xia et al., 2005; Kumar et al., 2006; Kumar et al., 2007; Brantley-Sieders et al., 2011).

The ligands to Eph receptors are called ephrins and they too are divided into two subfamilies – ephrin-A and ephrin-B – based on sequence homologies and interactions with Eph receptors (Arvanitis and Davy, 2012). In contrast to most other ligands of receptor tyrosine kinases, ephrins are also membrane-bound on the outside of cells – ephrin-A ligands *via* glycosphosphatidylinositol (GPI) links and ephrin-B with a single transmembrane domain.

Despite significant promiscuity between other Eph family members and the ephrin ligands, the single physiologically-relevant ligand of the EphB4 receptor is ephrin-B2 (Pasquale, 2004). In a similar manner to the classical trans-signalling pathways of Eph-ephrins, forward signalling, through EphB4, and reverse signalling, through ephrin-B2, requires heterotetramerisation of two receptors (on cell 1) and two ligands (on cell 2). This normal interaction between EphB4 and ephrin-B2, requires direct cell-cell contact and induces tumour suppression in the EphB4 expressing cell and stabilization of cell-cell adhesion and stimulation of angiogenesis through the ligand expressing cell (Chen et al., 2008; Pasquale, 2010). Disruption of cell-cell junctions or changes in the Eph-ephrin balance (through the overexpression of EphB4), both of which can occur in cancer cells, prevents or diminishes ligand-receptor interaction and allows the clustering of EphB4 and the use of ligand-independent signalling pathways that promote tumour formation. It has also been reported that Eph receptors and ephrins when co-expressed in neurons and cancer cells can engage in lateral cis associations, which lessen receptor activation through trans-signalling (Hornberger et al., 1999; Yin et al., 2004; Carvalho et al., 2006; Kao and Kania, 2011; Falivelli et al., 2013).

High affinity trans-signalling interactions between Eph receptors and ephrin ligands are terminated by two key mechanisms. In the first of these, the whole receptor-ligand complex is endocytosed into either cell *via* a mechanism that is dependent on Rac signalling and may include pinching off part of the plasma membrane of the opposite cell (Marston et al., 2003; Zimmer et al., 2003; Pitulescu and Adams, 2010). The second mechanism involves protease cleavage of the ligand or the EphR after formation of the receptor-ligand complex. Proteases linked to Eph receptor and ephrin proteins include RHBDL2, a rhomboid transmembrane serine protease that cleaves ephrin-B3, several matrix metalloproteinases and A Disintegrin and Metalloproteases (ADAMs) that have been reported to cleave both Eph receptors and ephrins, including ephrin-A2 and ephrin-B1 (Hattori et al., 2000; Pascall and Brown, 2004; Litterst et al., 2007; Tanaka et al., 2007; Lin et al., 2008; Inoue et al., 2009; Janes et al., 2009). Using indirect ‘gain of function’ approaches, EphB4 has been implicated as a potential substrate for ADAM-8, -9 and -17 (Janes et al., 2005; Guaiquil et al., 2009; Guaiquil et al., 2010; Mendelson et al., 2010; Weskamp et al., 2010).

More recently, we have identified EphB4 and its ephrin-B2 ligand as biochemical substrates of KLK4 (Lisle et al., 2013). An *in silico* approach was used to identify potential KLK4 cleavage sites through mapping the consensus KLK4 cleavage site to accessible epitopes on the surface

of proteins from protein databases, for which there were defined crystal structures. Ephrin-B2 was confirmed as a substrate of KLK4 by incubation of recombinant proteins in various molar ratios, and the primary cleavage site in ephrin-B2 confirmed using N-terminal sequencing as after R174 (NP_004084). Closely related kallikreins KLK3 (PSA), KLK2 and KLK4 can cleave ephrin-B2 and we have used a concentration of KLK peptidase at $>4 \mu\text{g/ml}$ that is under the physiological range as the PSA concentration is 100–2000 $\mu\text{g/ml}$ in normal male seminal fluid (Lovgren et al., 1999; Drabovich et al., 2011) while KLK2 is 2–20 $\mu\text{g/ml}$ (Lovgren et al., 1999) and KLK4 is 0.2–202 $\mu\text{g/ml}$ (Obiezu et al., 2005; Shaw and Diamandis, 2007).

Although EphB4 was not identified as a potential KLK4-substrate using the *in silico* approach (because a suitable crystal structure was not available), in parallel experiments recombinant EphB4 was also shown to be cleaved by recombinant KLK4 (Figure 2). The KLK4 cleavage site in EphB4 was identified by N-terminal sequencing and is in the extracellular domain between R508 and A509. Cleavage after R508 would release almost the complete extracellular domain of EphB4 and this was confirmed by addition of recombinant KLK4 to cells

expressing surface EphB4 with subsequent identification of the cleaved ectodomain fragment of the predicted size via Western blot analysis. In addition, a protein band of the predicted size of the cleaved ectodomain was identified in cell lysates from EphB4+/KLK4+ prostate cancer cells, confirming that cleavage of the ectodomain can occur in a biological setting. Interestingly, this cleavage event appears to precede a second cleavage event catalysed by the presenilin-dependent protease γ -secretase, which releases the intracellular part of EphB4 from the membrane. This mechanism – ectodomain shedding followed by release of an intracellular fragment, is common to many receptor tyrosine kinases and in many cases the released fragments have important biological functions (Stephenson et al., 2012). The biological consequences of these two EphB4 cleavage events, and the functions of the released fragments, are currently being explored in clinical prostate cancer samples and cell lines.

CUB-domain containing protein 1 (CDCP1)

In the last few years the cell surface glycoprotein CUB-domain containing protein 1 (CDCP1) has emerged as a novel substrate of serine proteinases that have Arg/Lys trypsin-like substrate specificity (He et al., 2010), including, potentially, those members of the KLK family that cleave after these amino acids, such as KLKs 1, 2, 4–6, 8, 10–14. However, it is not known if CDCP1 can be cleaved by those with chymotryptic-like specificity, i.e., KLKs 3, 7, 9, or with some cross chymotryptic substrate specificity. CDCP1, also known as SIMA135 (Hooper et al., 2003), gp140 (Brown et al., 2004), Trask (Bhatt et al., 2005) and CD318 (Wortmann et al., 2009), is an 836 amino acid single pass transmembrane glycoprotein that consists of extracellular and intracellular domains encompassing approximately 660 and 150 residues respectively. Serine proteinase-mediated cleavage occurs within the extracellular domain at adjacent sites, Arg368 and Lys369, to initiate Src-mediated phosphorylation of CDCP1 at tyrosine (Y) 734, Y743 and Y762 and binding of phospho(p)-Src and another kinase p-PKC δ (He et al., 2010). Recently it has been shown that serine proteinase-mediated cleavage of CDCP1 promotes the survival of disseminating cancer cells as they escape from blood vessels. Increased cancer cell dissemination occurs via a p-Src/p-PKC δ mechanism that initiates p-Akt pro-survival and anti-apoptotic pathways (Casar et al., 2012). CDCP1 is up-regulated in cancers of the lung, kidney, ovary and pancreas where its elevated




Figure 2 (A) Silver stained PAGE showing KLK4 cleavage of full length EphB4 extracellular domain tagged with IgG Fc tag (EphB4-ecd-Fc) at a ratio of active KLK4 to EphB4 of 1:10 (0.1), 1:100 (0.01) and 1:5,000 (0.005). The EphB4-ecd-Fc fragment generated by KLK4 activity is indicated by a red box. (B) Diagram showing KLK4 cleavage of human EphB4 releases an ectodomain (70 kDa) from the cell surface and leaves an intracellular membrane-tethered fragment (50 kDa). This cleavage is then followed by hydrolysis by γ -secretase releasing an intracellular domain (ICD) of 47 kDa.

expression is associated with poor patient outcomes (Awakura et al., 2008; Ikeda et al., 2009; Miyazawa et al., 2010; Razorenova et al., 2011; Dong et al., 2012). As CDCP1 is widely expressed in mammalian epithelia including these cancers of epithelial origin where elevated KLK expression is also commonly observed (Lawrence et al., 2010), we were interested to see if members of this family are able to cleave CDCP1. In our hands, low levels of KLK4 were able to induce CDCP1 cleavage within 6 h, generating a 70 kDa fragment (Figure 3A), with the schematic shown in Figure 3B, although the function of this cleaved product remains to be investigated. These data suggest that in settings where active KLK4 or other Arg/Lys-specific KLKs are present, CDCP1 may act as a plasma membrane substrate to transduce signals across the cell surface including that promoting cell survival.

CDCP1 is known to be cleaved by matriptase and trypsin, which also cleave PAR2 (Brown et al., 2004; He et al., 2010) but it is thought that the endogenous protease is plasmin as it was found to be necessary and sufficient to cleave CDCP1 *in vivo* and plasminogen knockout mice lacked cleavage of CDCP1 in lung tissue (Casar et al., 2012). Increasingly, the expression of protease activated receptors and other proteolytically-activated proteins such as CDCP1 is becoming more clinically significant as potential markers of disease progression and as molecular targets. For example up-regulation and aberrant expression of both CDCP1 and uPAR are correlated with a poorer prognosis and can be indicative of disease progression in some cancer types (Awakura et al., 2008; Ikeda et al., 2009; Ma and Tao, 2012). Furthermore, as plasmin is generated from plasminogen by uPA it is possible that CDCP1 and other protease activated receptors could be implicated in the same pathway, contributing to dissemination and metastasis of invading tumour cells and therefore could represent multiple targets within a common pathway. As we

noted for the HGF/Met activation and uPA/uPAR cascades, it appears likely that KLKs will also be able to contribute to activation of CDCP1 signalling at multiple levels as it has recently been shown that plasmin is the crucial serine protease executing *in vivo* cleavage of this receptor during cancer cell dissemination in mice (Casar et al., 2012).

Other cell membrane-bound proteins as KLK substrates

In addition to the above receptor systems, there are a range of other associated cell surface proteins that are cleaved by members of the KLK family. For example, KLK6 can hydrolyse amyloid precursor protein (APP), which is localized in senile plaques and neurofilament tangles in the brain of Alzheimer's disease (AD) patients and is important in the deposition of amyloid plaques during progression of this disease (Little et al., 1997; Ogawa et al., 2000; Magklara et al., 2003). Also, KLK6 is able to cleave myelin basic protein (MBP) in oligodendrocytes and Schwann cells in the central nerve system, which is related to demyelinating diseases such as multiple sclerosis (Scarlsbrick et al., 2012b). Although KLK5 and KLK7 have been localised in numerous tissues their function has been extensively elucidated in keratinization, stratum corneum formation, turnover and desquamation of the skin. This occurs via the hydrolysis and degradation of cell membrane-bound adhesion glycoproteins, such as corneodesmosin, desmocollin 1 and plakoglobin (Caubet et al., 2004).

The cadherin proteins, desmogleins (DSGs), are key factors in desmosome formation and cell integrity in particular in skin (Amagai and Stanley, 2012), and loss of them has also been associated with cancer progression (Thiery and Sleeman, 2006). DSG2 is a substrate of KLK7 and transfection of this peptidase significantly up-regulated levels of soluble DSG2 fragments in culture media from pancreatic cancer BxPC-3 cells (Ramani et al., 2008c). Interestingly, using siRNA knockdown and protease inhibition approaches, cleavage of DSG1 by KLK5 was induced in oral squamous cancer SCC25 cells where decreased aggregation of these cells was seen (Jiang et al., 2011). It is known that the cell-cell contact protein, E-cadherin, plays a critical role in cell integrity and loss or degradation of this molecule has been associated with cancer progression (Thiery and Sleeman, 2006). Of note, adding KLK7 into pancreatic cancer BxPC-3 cells generated more soluble E-cadherin fragments while the aggregation of these cells significantly decreased, but invasion increased (Johnson et al., 2007). These data suggest a role of these




Figure 3 (A) Anti-CDCP1 and -GAPDH Western blotting of lysates from HeLa-CDCP1 cells incubated with KLK4 (50 nM) at 37°C for 1, 6 and 24 h. Full-length (135 kDa) and cleaved (70 kDa) CDCP1 are indicated. (B) Schematic showing that KLK4 cleavage of full length CDCP1 generates a 70 kDa form.

KLK peptidases in the progression of cancer by hydrolysis of cell adhesion proteins.

Meprins are members of the astacin metalloprotease family and have α and β subunits originally identified on the apical brush or plasma membrane of epithelial cells (Bond et al., 2005). Their unique localization and proteomic activity have been associated with trafficking of host cells such as, macrophage or metastatic cancer cells (Bond et al., 2005). A recent study using a terminal amine isotopic labeling of substrates (TAILS) approach identified that pro-KLK7 is a substrate of meprin α and β in the conditioned media from keratinocyte HaCaT cells (Jefferson et al., 2012) supporting their function in skin epithelial cells. Interestingly, both meprin α and β subunits were localized in the plasma membrane of HaCaT keratinocytes and pro-meprin β was cleaved by KLK4 (Becker-Pauly et al., 2007; Ohler et al., 2010), further suggesting a complex interaction between these two protease families.

Conclusion

Here we have summarised the current knowledge on the hydrolysis of cell membrane receptors/proteins by KLK1 and other KLK peptidases (Table 1) and their initiation of downstream signalling pathways. Although there is a diversity of tissue specificity and expression of these proteases, we have identified a strong association between the classical membrane-bound receptor activation function of KLK1 and similar attributes of other more recently discovered members of the KLK family. In addition, it is notable that there is increasing evidence of complex proteolytic cascades between the KLKs, uPA/uPAR and metalloproteinase families. A better understanding of these interactive cascades and downstream cellular signalling pathways initiated by these peptidases will help to fully elucidate the biological functions and pathophysiological role of KLKs *in vivo*.

Acknowledgments: This work was supported by funds from the National Health and Medical Research Council (NHMRC) of Australia (grants 550523, 614206 JAC, YD, JDH), the Australia India Strategic Research Fund (grant BF060023, JAC, YD), The Cancer Council Queensland, Australian Prostate Cancer Foundation of Australia (JDH, JAC, YD, SAS), and Wesley Research Institute (JDH, JAC, YD). Queensland University of Technology (SAS, AH, JL), JDH is an Australian Research Council Future Fellow and JAC is a NHMRC Principal Research Fellow.

Disclosure of potential conflicts of interest: No potential conflicts of interest are disclosed.

References


- Abelous, J.E. and Bardier, E. (1909). Les substances hypotensives de l'urine humaine normale. *Compt. Rend. Soc. Biol.* 66, 511–512.
- Adams, M.N., Ramachandran, R., Yau, M.K., Suen, J.Y., Fairlie, D.P., Hollenberg, M.D., and Hooper, J.D. (2011). Structure, function and pathophysiology of protease activated receptors. *Pharmacol. Ther.* 130, 248–282.
- Almasi, C.E., Hoyer-Hansen, G., Christensen, I.J., Dano, K., and Pappot, H. (2005). Prognostic impact of liberated domain I of the urokinase plasminogen activator receptor in squamous cell lung cancer tissue. *Lung Cancer* 48, 349–355.
- Amagai, M. and Stanley, J.R. (2012). Desmoglein as a target in skin disease and beyond. *J. Invest. Dermatol.* 132, 776–784.
- Angelo, P.F., Lima, A.R., Alves, F.M., Blaber, S.I., Scarisbrick, I.A., Blaber, M., Juliano, L. and Juliano, M.A. (2006). Substrate specificity of human kallikrein 6: salt and glycosaminoglycan activation effects. *J. Biol. Chem.* 281, 3116–3126.
- Arvanitis, D.N. and Davy, A. (2012). Regulation and misregulation of Eph/ephrin expression. *Cell. Adh. Migr.* 6, 131–137.
- Awakura, Y., Nakamura, E., Takahashi, T., Kotani, H., Mikami, Y., Kadowaki, T., Myoumoto, A., Akiyama, H., Ito, N., Kamoto, T., et al. (2008). Microarray-based identification of CUB-domain containing protein 1 as a potential prognostic marker in conventional renal cell carcinoma. *J. Cancer Res. Clin. Oncol.* 134, 1363–1369.
- Beaufort, N., Debela, M., Creutzburg, S., Kellermann, J., Bode, W., Schmitt, M., Pidard, D., and Magdolen, V. (2006). Interplay of human tissue kallikrein 4 (hK4) with the plasminogen activation system: hK4 regulates the structure and functions of the urokinase-type plasminogen activator receptor (uPAR). *Biol. Chem.* 387, 217–222.
- Becker-Pauly, C., Howel, M., Walker, T., Vlad, A., Aufenvenne, K., Oji, V., Lottaz, D., Sterchi, E.E., Debela, M., Magdolen, V., et al. (2007). The alpha and beta subunits of the metalloprotease meprin are expressed in separate layers of human epidermis, revealing different functions in keratinocyte proliferation and differentiation. *J. Invest. Dermatol.* 127, 1115–1125.
- Bhatt, A.S., Erdjument-Bromage, H., Tempst, P., Craik, C.S., and Moasser, M.M. (2005). Adhesion signaling by a novel mitotic substrate of src kinases. *Oncogene* 24, 5333–5343.
- Bhoola, K.D., Figueroa, C.D., and Worthy, K. (1992). Bioregulation of kinins: kallikreins, kininogens, and kininases. *Pharmacol. Rev.* 44, 1–80.
- Biyashev, D., Tan, F., Chen, Z., Zhang, K., Deddish, P.A., Erdos, E.G., and Hecquet, C. (2006). Kallikrein activates bradykinin B2 receptors in absence of kininogen. *Am. J. Physiol. Heart Circ. Physiol.* 290, H1244–H1250.
- Bohm, S.K., Kong, W., Bromme, D., Smeekens, S.P., Anderson, D.C., Connolly, A., Kahn, M., Nelken, N.A., Coughlin, S.R., Payan, D.G., et al. (1996). Molecular cloning, expression and potential functions of the human proteinase-activated receptor-2. *Biochem. J.* 314, 1009–1016.
- Boire, A., Covic, L., Agarwal, A., Jacques, S., Sherifi, S., and Kuliopulos, A. (2005). PAR1 is a matrix metalloprotease-1

- receptor that promotes invasion and tumorigenesis of breast cancer cells. *Cell* 120, 303–313.
- Bond, J.S., Matters, G.L., Banerjee, S., and Dusheck, R.E. (2005). Meprin metalloprotease expression and regulation in kidney, intestine, urinary tract infections and cancer. *FEBS Lett.* 579, 3317–3322.
- Boyd, A.W., Bartlett, P.F., and Lackmann, M. (2014). Therapeutic targeting of EPH receptors and their ligands. *Nat. Rev. Drug Discov.* 13, 39–62.
- Brady, J.M., Wines, D.R., and MacDonald, R.J. (1989). Expression of two kallikrein gene family members in the rat prostate. *Biochemistry* 28, 5203–5210.
- Brantley-Sieders, D.M., Jiang, A., Sarma, K., Badu-Nkansah, A., Walter, D.L., Shyr, Y., and Chen, J. (2011). Eph/ephrin profiling in human breast cancer reveals significant associations between expression level and clinical outcome. *PLoS One* 6, e24426.
- Brown, T.A., Yang, T.M., Zaitsevskaya, T., Xia, Y., Dunn, C.A., Sigle, R.O., Knudsen, B., and Carter, W.G. (2004). Adhesion or plasmin regulates tyrosine phosphorylation of a novel membrane glycoprotein p80/gp140/CUB domain-containing protein 1 in epithelia. *J. Biol. Chem.* 279, 14772–14783.
- Bush, J.O. and Soriano, P. (2012). Eph/ephrin signaling: genetic, phosphoproteomic, and transcriptomic approaches. *Semin. Cell Dev. Biol.* 23, 26–34.
- Carvalho, R.F., Beutler, M., Marler, K.J., Knoll, B., Becker-Barroso, E., Heintzmann, R., Ng, T., and Drescher, U. (2006). Silencing of EphA3 through a cis interaction with ephrinA5. *Nat. Neurosci.* 9, 322–330.
- Casar, B., He, Y., Ionomou, M., Hooper, J.D., Quigley, J.P., and Der-yugina, E.I. (2012). Blocking of CDCP1 cleavage in vivo prevents Akt-dependent survival and inhibits metastatic colonization through PARP1-mediated apoptosis of cancer cells. *Oncogene* 31, 3924–3938.
- Caubet, C., Jonca, N., Brattsand, M., Guerrin, M., Bernard, D., Schmidt, R., Egelrud, T., Simon, M., and Serre, G. (2004). Degradation of corneodesmosome proteins by two serine proteases of the kallikrein family, SCTE/KLK5/hK5 and SCCE/KLK7/hK7. *J. Invest. Dermatol.* 122, 1235–1244.
- Chao, J., Buse, J., Shimamoto, K., and Margolius, H.S. (1981). Kallikrein-induced uterine contraction independent of kinin formation. *Proc. Natl. Acad. Sci. USA* 78, 6154–6157.
- Chao, J., Bledsoe, G., Yin, H., and Chao, L. (2006). The tissue kallikrein-kinin system protects against cardiovascular and renal diseases and ischemic stroke independently of blood pressure reduction. *Biol. Chem.* 387, 665–675.
- Chao, J., Yin, H., Gao, L., Hagiwara, M., Shen, B., Yang, Z.R., and Chao, L. (2008). Tissue kallikrein elicits cardioprotection by direct kinin b2 receptor activation independent of kinin formation. *Hypertension* 52, 715–720.
- Chen, J., Zhuang, G., Frieden, L., and Debinski, W. (2008). Eph receptors and Ephrins in cancer: common themes and controversies. *Cancer Res.* 68, 10031–10033.
- Chung, H., Hamza, M., Oikonomopoulou, K., Gratio, V., Saifedine, M., Virca, G.D., Diamandis, E.P., Hollenberg, M.D., and Darmoul, D. (2012). Kallikrein-related peptidase signaling in colon carcinoma cells: targeting proteinase-activated receptors (PARs). *Biol. Chem.* 393, 413–420.
- Clements, J.A. (2000). Tissue kallikrein–kinin system. In: *Handbook of Physiology: The Endocrine System, Endocrine Regulation of Water and Electrolyte Balance*, J.C.S. Fray, ed. (New York: Oxford University Press), pp. 333–376.
- da Costa, P.L., Sirois, P., Tannock, I.F., and Chammas, R. (2014). The role of kinin receptors in cancer and therapeutic opportunities. *Cancer Lett.* 345, 27–38.
- Dong, Y., He, Y., de Boer, L., Stack, M.S., Lumley, J.W., Clements, J.A., and Hooper, J.D. (2012). The cell surface glycoprotein CUB domain-containing protein 1 (CDCP1) contributes to epidermal growth factor receptor-mediated cell migration. *J. Biol. Chem.* 287, 9792–9803.
- Drabovich, A.P., Jarvi, K., and Diamandis, E.P. (2011). Verification of male infertility biomarkers in seminal plasma by multiplex selected reaction monitoring assay. *Mol. Cell. Proteomics* 10, M110 004127.
- Falivelli, G., Lisabeth, E.M., Rubio de la Torre, E., Perez-Tenorio, G., Tosato, G., Salvucci, O., and Pasquale, E.B. (2013). Attenuation of eph receptor kinase activation in cancer cells by coexpressed ephrin ligands. *PLoS One* 8, e81445.
- Fiedler, F. (1979). Enzymology of glandular kallikreins. In: *Bradykinin, kallidin, and kallikrein*. E.G. Erdos, ed. (Berlin, Heidelberg, New York: Springer-Verlag), pp. 101–161.
- Frenette, G., Tremblay, R.R., Lazure, C. and Dube, J.Y. (1997). Prostatic kallikrein hK2, but not prostate-specific antigen (hK3), activates single-chain urokinase-type plasminogen activator. *Int. J. Cancer* 71, 897–899.
- Gan, L., Lee, I., Smith, R., Argonza-Barrett, R., Lei, H., McCuaig, J., Moss, P., Paeper, B. and Wang, K. (2000). Sequencing and expression analysis of the serine protease gene cluster located in chromosome 19q13 region. *Gene* 257, 119–130.
- Gao, L., Chao, L. and Chao, J. (2010a). A novel signaling pathway of tissue kallikrein in promoting keratinocyte migration: activation of proteinase-activated receptor 1 and epidermal growth factor receptor. *Exp. Cell Res.* 316, 376–389.
- Gao, L., Smith, R.S., Chen, L.M., Chai, K.X., Chao, L. and Chao, J. (2010b). Tissue kallikrein promotes prostate cancer cell migration and invasion via a protease-activated receptor-1-dependent signaling pathway. *Biol. Chem.* 391, 803–812.
- Gratio, V., Beaufort, N., Seiz, L., Maier, J., Virca, G.D., Debela, M., Grebenchtchikov, N., Magdolen, V. and Darmoul, D. (2010). Kallikrein-related peptidase 4: a new activator of the aberrantly expressed protease-activated receptor 1 in colon cancer cells. *Am. J. Pathol.* 176, 1452–1461.
- Gratio, V., Lorient, C., Virca, G.D., Oikonomopoulou, K., Walker, F., Diamandis, E.P., Hollenberg, M.D. and Darmoul, D. (2011). Kallikrein-related peptidase 14 acts on proteinase-activated receptor 2 to induce signaling pathway in colon cancer cells. *Am. J. Pathol.* 179, 2625–2636.
- Guaiquil, V., Swendeman, S., Yoshida, T., Chavala, S., Campochiaro, P.A. and Blobel, C.P. (2009). ADAM9 is involved in pathological retinal neovascularization. *Mol. Cell. Biol.* 29, 2694–2703.
- Guaiquil, V.H., Swendeman, S., Zhou, W., Guaiquil, P., Weskamp, G., Bartsch, J.W. and Blobel, C.P. (2010). ADAM8 is a negative regulator of retinal neovascularization and of the growth of heterotopically injected tumor cells in mice. *J. Mol. Med. (Berl.)* 88, 497–505.
- Harvey, T.J., Hooper, J.D., Myers, S.A., Stephenson, S.A., Ashworth, L.K., and Clements, J.A. (2000). Tissue-specific expression patterns and fine mapping of the human kallikrein (KLK) locus on proximal 19q13.4. *J. Biol. Chem.* 275, 37397–37406.
- Hattori, M., Osterfield, M., and Flanagan, J.G. (2000). Regulated cleavage of a contact-mediated axon repellent. *Science* 289, 1360–1365.

- He, Y., Wortmann, A., Burke, L.J., Reid, J.C., Adams, M.N., Abdul-Jabar, I., Quigley, J.P., Leduc, R., Kirchhofer, D., and Hooper, J.D. (2010). Proteolysis-induced N-terminal ectodomain shedding of the integral membrane glycoprotein CUB domain-containing protein 1 (CDCP1) is accompanied by tyrosine phosphorylation of its C-terminal domain and recruitment of Src and PKCdelta. *J. Biol. Chem.* 285, 26162–26173.
- Hecquet, C., Tan, F., Marcic, B.M., and Erdos, E.G. (2000). Human bradykinin B(2) receptor is activated by kallikrein and other serine proteases. *Mol. Pharmacol.* 58, 828–836.
- Hooper, J.D., Zijlstra, A., Aimes, R.T., Liang, H., Claassen, G.F., Tarin, D., Testa, J.E., and Quigley, J.P. (2003). Subtractive immunization using highly metastatic human tumor cells identifies SIMA135/CDCP1, a 135 kDa cell surface phosphorylated glycoprotein antigen. *Oncogene* 22, 1783–1794.
- Hornberger, M.R., Dutting, D., Ciossek, T., Yamada, T., Handwerker, C., Lang, S., Weth, F., Huf, J., Wessel, R., Logan, C., et al. (1999). Modulation of EphA receptor function by coexpressed ephrinA ligands on retinal ganglion cell axons. *Neuron* 22, 731–742.
- Huai, Q., Mazar, A.P., Kuo, A., Parry, G.C., Shaw, D.E., Callahan, J., Li, Y., Yuan, C., Bian, C., Chen, L., et al. (2006). Structure of human urokinase plasminogen activator in complex with its receptor. *Science*. 311, 656–659.
- Ikeda, J., Oda, T., Inoue, M., Uekita, T., Sakai, R., Okumura, M., Aozasa, K., and Morii, E. (2009). Expression of CUB domain containing protein (CDCP1) is correlated with prognosis and survival of patients with adenocarcinoma of lung. *Cancer Sci.* 100, 429–433.
- Inoue, E., Deguchi-Tawarada, M., Togawa, A., Matsui, C., Arita, K., Katahira-Tayama, S., Sato, T., Yamauchi, E., Oda, Y., and Takai, Y. (2009). Synaptic activity prompts gamma-secretase-mediated cleavage of EphA4 and dendritic spine formation. *J. Cell Biol.* 185, 551–564.
- Ishihara, H., Connolly, A.J., Zeng, D., Kahn, M.L., Zheng, Y.W., Timmons, C., Tram, T., and Coughlin, S.R. (1997). Protease-activated receptor 3 is a second thrombin receptor in humans. *Nature* 386, 502–506.
- Janes, P.W., Saha, N., Barton, W.A., Kolev, M.V., Wimmer-Kleikamp, S.H., Nievergall, E., Blobel, C.P., Himanen, J.P., Lackmann, M., and Nikolov, D.B. (2005). Adam meets Eph: an ADAM substrate recognition module acts as a molecular switch for ephrin cleavage in trans. *Cell* 123, 291–304.
- Janes, P.W., Wimmer-Kleikamp, S.H., Frangakis, A.S., Treble, K., Griesshaber, B., Sabet, O., Grabenbauer, M., Ting, A.Y., Saftig, P., Bastiaens, P.I., et al. (2009). Cytoplasmic relaxation of active Eph controls ephrin shedding by ADAM10. *PLoS Biol.* 7, e1000215.
- Jefferson, T., Auf dem Keller, U., Bellac, C., Metz, V.V., Broder, C., Hedrich, J., Ohler, A., Maier, W., Magdolen, V., Sterchi, E., et al. (2012). The substrate degradome of meprin metalloproteases reveals an unexpected proteolytic link between meprin beta and ADAM10. *Cell. Mol. Life Sci.* 70, 309–333.
- Jiang, R., Shi, Z., Johnson, J.J., Liu, Y., and Stack, M.S. (2011). Kallikrein-5 promotes cleavage of desmoglein-1 and loss of cell-cell cohesion in oral squamous cell carcinoma. *J. Biol. Chem.* 286, 9127–9135.
- Johnson, S.K., Ramani, V.C., Hennings, L., and Haun, R.S. (2007). Kallikrein 7 enhances pancreatic cancer cell invasion by shedding E-cadherin. *Cancer* 109, 1811–1820.
- Kao, T.J. and Kania, A. (2011). Ephrin-mediated cis-attenuation of Eph receptor signaling is essential for spinal motor axon guidance. *Neuron*. 71, 76–91.
- Kayashima, Y., Smithies, O. and Kakoki, M. (2012). The kallikrein-kinin system and oxidative stress. *Curr. Opin. Nephrol. Hypertens.* 21, 92–96.
- Kraut, H., Frey, E.K., and Werle, E. (1930). Der nachweis eines kreislauf-hormons in der pankreasdrüse. *Hoppe-Seyler's Z. Physiol. Chem.* 189, 97–106.
- Kumar, S.R., Singh, J., Xia, G., Krasnoperov, V., Hassanieh, L., Ley, E.J., Scheinet, J., Kumar, N.G., Hawes, D., Press, M.F., et al. (2006). Receptor tyrosine kinase EphB4 is a survival factor in breast cancer. *Am. J. Pathol.* 169, 279–293.
- Kumar, S.R., Masood, R., Spannuth, W.A., Singh, J., Scheinet, J., Kleiber, G., Jennings, N., Deavers, M., Krasnoperov, V., Dubeau, L., et al. (2007). The receptor tyrosine kinase EphB4 is overexpressed in ovarian cancer, provides survival signals and predicts poor outcome. *Br. J. Cancer* 96, 1083–1091.
- Lawrence, M.G., Lai, J., and Clements, J.A. (2010). Kallikreins on steroids: structure, function, and hormonal regulation of prostate-specific antigen and the extended kallikrein locus. *Endocr. Rev.* 31, 407–446.
- Lee, Y.C., Perren, J.R., Douglas, E.L., Raynor, M.P., Bartley, M.A., Bardy, P.G., and Stephenson, S.A. (2005). Investigation of the expression of the EphB4 receptor tyrosine kinase in prostate carcinoma. *B.M.C. Cancer* 5, 119.
- Lin, K.T., Sloniowski, S., Ethell, D.W., and Ethell, I.M. (2008). Ephrin-B2-induced cleavage of EphB2 receptor is mediated by matrix metalloproteinases to trigger cell repulsion. *J. Biol. Chem.* 283, 28969–28979.
- Lisle, J.E., Mertens-Walker, I., Rutkowski, R., Herington, A.C., and Stephenson, S.A. (2013). Eph receptors and their ligands: promising molecular biomarkers and therapeutic targets in prostate cancer. *Biochim. Biophys. Acta* 1835, 243–257.
- Litterst, C., Georgakopoulos, A., Shioi, J., Ghersi, E., Wisniewski, T., Wang, R., Ludwig, A., and Robakis, N.K. (2007). Ligand binding and calcium influx induce distinct ectodomain/gamma-secretase-processing pathways of EphB2 receptor. *J. Biol. Chem.* 282, 16155–16163.
- Little, S.P., Dixon, E.P., Norris, F., Buckley, W., Becker, G.W., Johnson, M., Dobbins, J.R., Wyrick, T., Miller, J.R., MacKellar, W., et al. (1997). Zyme, a novel and potentially amyloidogenic enzyme cDNA isolated from Alzheimer's disease brain. *J. Biol. Chem.* 272, 25135–25142.
- Lovgren, J., Valtonen-Andre, C., Marsal, K., Lilja, H., and Lundwall, A. (1999). Measurement of prostate-specific antigen and human glandular kallikrein 2 in different body fluids. *J. Androl.* 20, 348–355.
- Ma, Y.Y. and Tao, H.Q. (2012). Role of urokinase plasminogen activator receptor in gastric cancer: a potential therapeutic target. *Cancer Biother. Radiopharm.* 27, 285–290.
- Macfarlane, S.R., Seatter, M.J., Kanke, T., Hunter, G.D., and Plevin, R. (2001). Proteinase-activated receptors. *Pharmacol. Rev.* 53, 245–282.
- Madeddu, P., Emanuelli, C., and El-Dahr, S. (2007). Mechanisms of disease: the tissue kallikrein-kinin system in hypertension and vascular remodeling. *Nat. Clin. Pract. Nephrol.* 3, 208–221.
- Magklara, A., Mellati, A.A., Wasney, G.A., Little, S.P., Sotiropoulou, G., Becker, G.W., and Diamandis, E.P. (2003). Characterization of the enzymatic activity of human kallikrein 6: Autoactivation,

- substrate specificity, and regulation by inhibitors. *Biochem. Biophys. Res. Commun.* 307, 948–955.
- Mahtouk, K., Tjin, E.P., Spaargaren, M., and Pals, S.T. (2010). The HGF/MET pathway as target for the treatment of multiple myeloma and B-cell lymphomas. *Biochim. Biophys. Acta.* 1806, 208–219.
- Marcondes, S. and Antunes, E. (2005). The plasma and tissue kininogen-kallikrein-kinin system: role in the cardiovascular system. *Curr. Med. Chem. Cardiovasc. Hematol. Agents* 3, 33–44.
- Mars, W.M., Zarnegar, R., and Michalopoulos, G.K. (1993). Activation of hepatocyte growth factor by the plasminogen activators uPA and tPA. *Am. J. Pathol.* 143, 949–958.
- Marston, D.J., Dickinson, S., and Nobes, C.D. (2003). Rac-dependent trans-endocytosis of ephrinBs regulates Eph-ephrin contact repulsion. *Nat. Cell Biol.* 5, 879–888.
- Mendelson, K., Swendeman, S., Saftig, P., and Blobel, C.P. (2010). Stimulation of platelet-derived growth factor receptor beta (PDGFRbeta) activates ADAM17 and promotes metalloproteinase-dependent cross-talk between the PDGFRbeta and epidermal growth factor receptor (EGFR) signaling pathways. *J. Biol. Chem.* 285, 25024–25032.
- Miyazawa, Y., Uekita, T., Hiraoka, N., Fujii, S., Kosuge, T., Kanai, Y., Nojima, Y., and Sakai, R. (2010). CUB domain-containing protein 1, a prognostic factor for human pancreatic cancers, promotes cell migration and extracellular matrix degradation. *Cancer Res.* 70, 5136–5146.
- Mize, G.J., Wang, W., and Takayama, T.K. (2008). Prostate-specific kallikreins-2 and -4 enhance the proliferation of DU-145 prostate cancer cells through protease-activated receptors-1 and -2. *Mol. Cancer Res.* 6, 1043–1051.
- Mukai, S., Fukushima, T., Naka, D., Tanaka, H., Osada, Y., and Kataoka, H. (2008). Activation of hepatocyte growth factor activator zymogen (pro-HGFA) by human kallikrein 1-related peptidases. *FEBS J.* 275, 1003–1017.
- Naldini, L., Tamagnone, L., Vigna, E., Sachs, M., Hartmann, G., Birchmeier, W., Daikuhara, Y., Tsubouchi, H., Blasi, F., and Comoglio, P.M. (1992). Extracellular proteolytic cleavage by urokinase is required for activation of hepatocyte growth factor/scatter factor. *EMBO J.* 11, 4825–4833.
- Nievergall, E., Saunders, T., and Lackmann, M. (2012). Targeting of EPH receptor tyrosine kinases for anticancer therapy. *Crit. Rev. Oncog.* 17, 211–232.
- Obiezu, C.V., Shan, S.J., Soosaipillai, A., Luo, L.Y., Grass, L., Sotiropoulou, G., Petraki, C.D., Papanastasiou, P.A., Levesque, M.A., and Diamandis, E.P. (2005). Human kallikrein 4: quantitative study in tissues and evidence for its secretion into biological fluids. *Clin. Chem.* 51, 1432–1442.
- Ogawa, K., Yamada, T., Tsujioka, Y., Taguchi, J., Takahashi, M., Tsuboi, Y., Fujino, Y., Nakajima, M., Yamamoto, T., Akatsu, H., et al. (2000). Localization of a novel type trypsin-like serine protease, neurosin, in brain tissues of Alzheimer's disease and Parkinson's disease. *Psychiatry Clin. Neurosci.* 54, 419–426.
- Ohler, A., Debela, M., Wagner, S., Magdolen, V., and Becker-Paul, C. (2010). Analyzing the protease web in skin: meprin metalloproteases are activated specifically by KLK4, 5 and 8 vice versa leading to processing of proKLK7 thereby triggering its activation. *Biol. Chem.* 391, 455–460.
- Oikonomopoulou, K., Hansen, K.K., Saifeddine, M., Tea, I., Blaber, M., Blaber, S.I., Scarisbrick, I., Andrade-Gordon, P., Cottrell, G.S., Bunnett, N.W., et al. (2006). Proteinase-activated receptors, targets for kallikrein signaling. *J. Biol. Chem.* 281, 32095–32112.
- Pascall, J.C. and Brown, K.D. (2004). Intramembrane cleavage of ephrinB3 by the human rhomboid family protease, RHBDL2. *Biochem. Biophys. Res. Commun.* 317, 244–252.
- Pasquale, E.B. (2004). Eph-ephrin promiscuity is now crystal clear. *Nat. Neurosci.* 7, 417–418.
- Pasquale, E.B. (2010). Eph receptors and ephrins in cancer: bidirectional signalling and beyond. *Nat. Rev. Cancer.* 10, 165–180.
- Pitulescu, M.E. and Adams, R.H. (2010). Eph/ephrin molecules--a hub for signaling and endocytosis. *Genes Dev.* 24, 2480–2492.
- Rajapakse, S., Ogiwara, K., Takano, N., Moriyama, A., and Takahashi, T. (2005). Biochemical characterization of human kallikrein 8 and its possible involvement in the degradation of extracellular matrix proteins. *FEBS Lett.* 579, 6879–6884.
- Ramachandran, R., Eissa, A., Mihara, K., Oikonomopoulou, K., Saifeddine, M., Renaux, B., Diamandis, E., and Hollenberg, M.D. (2012). Proteinase-activated receptors (PARs): differential signalling by kallikrein-related peptidases KLK8 and KLK14. *Biol. Chem.* 393, 421–427.
- Ramani, V.C. and Haun, R.S. (2008a). Expression of kallikrein 7 diminishes pancreatic cancer cell adhesion to vitronectin and enhances urokinase-type plasminogen activator receptor shedding. *Pancreas.* 37, 399–404.
- Ramani, V.C. and Haun, R.S. (2008b). The extracellular matrix protein fibronectin is a substrate for kallikrein 7. *Biochem. Biophys. Res. Commun.* 369, 1169–1173.
- Ramani, V.C., Hennings, L., and Haun, R.S. (2008c). Desmoglein 2 is a substrate of kallikrein 7 in pancreatic cancer. *BMC Cancer* 8, 373.
- Ramsay, A.J., Dong, Y., Hunt, M.L., Linn, M., Samarasinghe, H., Clements, J.A. and Hooper, J.D. (2008). Kallikrein-related peptidase 4 (KLK4) initiates intracellular signaling via protease-activated receptors (PARs). KLK4 and PAR-2 are co-expressed during prostate cancer progression. *J. Biol. Chem.* 283, 12293–12304.
- Rasch, M.G., Lund, I.K., Almasi, C.E., and Hoyer-Hansen, G. (2008). Intact and cleaved uPAR forms: diagnostic and prognostic value in cancer. *Front. Biosci.* 13, 6752–6762.
- Razorenova, O.V., Finger, E.C., Colavitti, R., Chernikova, S.B., Boiko, A.D., Chan, C.K., Krieg, A., Bedogni, B., LaGory, E., Weissman, I.L., et al. (2011). VHL loss in renal cell carcinoma leads to up-regulation of CUB domain-containing protein 1 to stimulate PKC{delta}-driven migration. *Proc. Natl. Acad. Sci. USA* 108, 1931–1936.
- Regoli, D., Plante, G.E. and Gobeil, F., Jr. (2012). Impact of kinins in the treatment of cardiovascular diseases. *Pharmacol. Ther.* 135, 94–111.
- Reuning, U., Sperl, S., Kopitz, C., Kessler, H., Kruger, A., Schmitt, M., and Magdolen, V. (2003). Urokinase-type plasminogen activator (uPA) and its receptor (uPAR): development of antagonists of uPA/uPAR interaction and their effects *in vitro* and *in vivo*. *Curr. Pharm. Des.* 9, 1529–1543.
- Rhaleb, N.E., Yang, X.P., and Carretero, O.A. (2011). The kallikrein-kinin system as a regulator of cardiovascular and renal function. *Compr. Physiol.* 1, 971–993.
- Rocha e Silva, M., Beraldo, W.T., and Rosenfeld, G. (1949). Bradykinin, a hypotensive and smooth muscle stimulating factor released from plasma globulin by snake venoms and by trypsin. *Am. J. Physiol.* 156, 261–273.

- Savvatis, K., Westermann, D., Schultheiss, H.P., and Tschope, C. (2010). Kinins in cardiac inflammation and regeneration: insights from ischemic and diabetic cardiomyopathy. *Neuropeptides* 44, 119–125.
- Scarlsbrick, I.A., Epstein, B., Cloud, B.A., Yoon, H., Wu, J., Renner, D.N., Blaber, S.I., Blaber, M., Vandell, A.G., and Bryson, A.L. (2011). Functional role of kallikrein 6 in regulating immune cell survival. *PLoS One* 6, e18376.
- Scarlsbrick, I.A., Radulovic, M., Burda, J.E., Larson, N., Blaber, S.I., Giannini, C., Blaber, M., and Vandell, A.G. (2012a). Kallikrein 6 is a novel molecular trigger of reactive astrogliosis. *Biol. Chem.* 393, 355–367.
- Scarlsbrick, I.A., Yoon, H., Panos, M., Larson, N., Blaber, S.I., Blaber, M., and Rodriguez, M. (2012b). Kallikrein 6 regulates early CNS demyelination in a viral model of multiple sclerosis. *Brain Pathol.* 22, 709–722.
- Schachter, M. (1979). Kallikreins (kininogenases)--a group of serine proteases with bioregulatory actions. *Pharmacol. Rev.* 31, 1–17.
- Schmalfeldt, B., Prechtel, D., Harting, K., Spathe, K., Rutke, S., Konik, E., Fridman, R., Berger, U., Schmitt, M., Kuhn, W., et al. (2001). Increased expression of matrix metalloproteinases (MMP)-2, MMP-9, and the urokinase-type plasminogen activator is associated with progression from benign to advanced ovarian cancer. *Clin. Cancer Res.* 7, 2396–2404.
- Shaw, J.L. and Diamandis, E.P. (2007). Distribution of 15 human kallikreins in tissues and biological fluids. *Clin. Chem.* 53, 1423–1432.
- Smith, H.W. and Marshall, C.J. (2010). Regulation of cell signalling by uPAR. *Nat Rev Mol. Cell. Biol.* 11, 23–36.
- Spinetti, G., Fortunato, O., Cordella, D., Portararo, P., Krankel, N., Katare, R., Sala-Newby, G.B., Richer, C., Vincent, M.P., Alhenc-Gelas, F., et al. (2011). Tissue kallikrein is essential for invasive capacity of circulating proangiogenic cells. *Circ. Res.* 108, 284–293.
- Stadnicki, A. (2011). Intestinal tissue kallikrein-kinin system in inflammatory bowel disease. *Inflamm. Bowel Dis.* 17, 645–654.
- Stefansson, K., Brattsand, M., Roosterman, D., Kempkes, C., Bocheva, G., Steinhoff, M., and Egelrud, T. (2008). Activation of proteinase-activated receptor-2 by human kallikrein-related peptidases. *J. Invest. Dermatol.* 128, 18–25.
- Stephenson, S.A., Mertens-Walker, I., and Herington, A.C. (2012). Signaling of receptor tyrosine kinases in the nucleus. In: *Current Frontiers and Perspectives in Cell Biology*, S.R. Najman, ed. (Croatia: InTech), pp. 211–234.
- Takayama, T.K., McMullen, B.A., Nelson, P.S., Matsumura, M., and Fujikawa, K. (2001). Characterization of hK4 (prostate), a prostate-specific serine protease: activation of the precursor of prostate specific antigen (pro-PSA) and single-chain urokinase-type plasminogen activator and degradation of prostatic acid phosphatase. *Biochemistry* 40, 15341–15348.
- Tanaka, M., Sasaki, K., Kamata, R., and Sakai, R. (2007). The C-terminus of ephrin-B1 regulates metalloproteinase secretion and invasion of cancer cells. *J. Cell Sci.* 120, 2179–2189.
- Thiery, J.P. and Sleeman, J.P. (2006). Complex networks orchestrate epithelial-mesenchymal transitions. *Nat. Rev. Mol. Cell. Biol.* 7, 131–142.
- Trusolino, L., Bertotti, A., and Comoglio, P.M. (2010). MET signalling: principles and functions in development, organ regeneration and cancer. *Nat. Rev. Mol. Cell. Biol.* 11, 834–848.
- Vandell, A.G., Larson, N., Laxmikanthan, G., Panos, M., Blaber, S.I., Blaber, M., and Scarlsbrick, I.A. (2008). Protease-activated receptor dependent and independent signaling by kallikreins 1 and 6 in CNS neuron and astroglial cell lines. *J. Neurochem.* 107, 855–870.
- Vesey, D.A., Hooper, J.D., Gobe, G.C., and Johnson, D.W. (2007). Potential physiological and pathophysiological roles for protease-activated receptor-2 in the kidney. *Nephrology (Carlton)* 12, 36–43.
- Viel, T.A. and Buck, H.S. (2011). Kallikrein-kinin system mediated inflammation in Alzheimer's disease *in vivo*. *Curr. Alzheimer Res.* 8, 59–66.
- Vu, T.K., Hung, D.T., Wheaton, V.I., and Coughlin, S.R. (1991). Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell* 64, 1057–1068.
- Wang, W., Mize, G.J., Zhang, X., and Takayama, T.K. (2009). Kallikrein-related peptidase-4 initiates tumor-stroma interactions in prostate cancer through protease-activated receptor-1. *Int. J. Cancer.* 126, 599–610.
- Werle, E. (1934). Zur Kenntnis des haushalts des Kallikreins. *Biochem Z.* 269, 415–434.
- Weskamp, G., Mendelson, K., Swendeman, S., Le Gall, S., Ma, Y., Lyman, S., Hinoki, A., Eguchi, S., Guaiquil, V., Horiuchi, K., et al. (2010). Pathological neovascularization is reduced by inactivation of ADAM17 in endothelial cells but not in pericytes. *Circ. Res.* 106, 932–940.
- Wortmann, A., He, Y., Deryugina, E.I., Quigley, J.P., and Hooper, J.D. (2009). The cell surface glycoprotein CD133 in cancer--insights, opportunities, and challenges. *IUBMB Life* 61, 723–730.
- Xia, G., Kumar, S.R., Masood, R., Zhu, S., Reddy, R., Krasnoperov, V., Quinn, D.I., Henshall, S.M., Sutherland, R.L., Pinski, J.K., et al. (2005). EphB4 expression and biological significance in prostate cancer. *Cancer Res.* 65, 4623–4632.
- Xu, W.F., Andersen, H., Whitmore, T.E., Presnell, S.R., Yee, D.P., Ching, A., Gilbert, T., Davie, E.W., and Foster, D.C. (1998). Cloning and characterization of human protease-activated receptor 4. *Proc. Natl. Acad. Sci. USA* 95, 6642–6646.
- Yin, Y., Yamashita, Y., Noda, H., Okafuji, T., Go, M.J., and Tanaka, H. (2004). EphA receptor tyrosine kinases interact with co-expressed ephrin-A ligands in cis. *Neurosci. Res.* 48, 285–296.
- Yousef, G.M. and Diamandis, E.P. (2001). The new human tissue kallikrein gene family: structure, function, and association to disease. *Endocr. Rev.* 22, 184–204.
- Zimmer, M., Palmer, A., Kohler, J., and Klein, R. (2003). EphB-ephrinB bi-directional endocytosis terminates adhesion allowing contact mediated repulsion. *Nat. Cell. Biol.* 5, 869–878.




Ying Dong studied Medicine at Beijing Union Medical College, China, where she then worked as an obstetrician and gynaecologist. Her PhD study was on biomarkers for ovarian cancer progression and treatment responses at the University of Queensland. She is a Research Fellow in the Cancer Program, Translational Research Institute, Queensland University of Technology. Her research is focused on understanding the functions of kallikrein proteases and the role of the tumour microenvironment in ovarian cancer.



Brittney Harrington completed her studies of Bachelor of Science with Honours at the University of Queensland. Brittney is currently performing research for her PhD at the Mater Medical Research Institute studying the role of the cell membrane protein CDCP1 in ovarian cancer progression, in particular in association with the tumour microenvironment.



Mark Adams undertook his undergraduate and honours degree studies at the Queensland University of Technology. He completed his PhD in 2012 at the Mater Medical Research Institute whilst supervised by Associate Professor John Hooper studying the protease-activated receptor-2 in cancer. He currently works as a postdoctoral research fellow based at the Translational Research Institute, Queensland University of Technology, examining the role of genomic instability in cancer progression.



Andreas Wortmann studied Biotechnology at the University of Applied Sciences Furtwangen in Germany. After three years in the

pharmaceutical industry working on the pre-clinical development of anti-tumour compounds, he studied for his PhD in cancer cell biology at the Institute of Health and Biomedical Innovation, Queensland University of Technology. In early 2011, he received his PhD and continued with two years of post-doctoral research in cancer cell biology and medical bionics. Since early 2013, he has worked at the Australian Therapeutic Goods Administration, in the evaluation of risk-management plans for high risk prescription medicines.



Sally-Anne Stephenson completed a PhD at the University of Queensland, Australia in 1998. Following post-doctoral studies at Queensland University of Technology (QUT), Brisbane where she identified KLK4 in the human kallikrein locus and at The Queen Elizabeth Hospital, Adelaide where she identified EphB4 as a protein increased in colon cancer, she started the Eph receptor biology group now at QUT's Institute of Health and Biomedical Innovation and part of the Translational Research Institute.



Jessica Lisle gained both her Bachelor's degree and Honours in Applied Science from the Queensland University of Technology. She is currently a PhD student studying protease regulation of the EphB4 receptor by kallikrein-related peptidase 4 in prostate cancer at the Translational Research Institute, Queensland University of Technology.



Professor Adrian Herington studied Biochemistry at Monash University in Melbourne, Australia, where he received his PhD for studies in the growth hormone (GH) and insulin-like growth factor (IGF) fields in 1973. After postdoctoral work in the USA, he continued this work leading independent research groups at the Prince Henry's Hospital Research Centre and the Royal Children's Hospital in Melbourne, until moving to the Queensland University of Technology in Brisbane in 1995. Currently, he heads QUT's Institute of Health & Biomedical Innovation's research teams at the Translational Research Institute in Brisbane. His personal research focusses on both the ghrelin axis and the receptor tyrosine kinase EphB4 in hormone-dependent cancers of the prostate and breast.



John Hooper studied chemistry at the University of Queensland and performed his PhD research on serine proteases in cancer and other pathologies at the Queensland Institute of Medical Research. His postdoctoral work at The Scripps Research Institute involved the use of animal models to examine proteases and receptor systems in metastasis and angiogenesis. His laboratory at the Mater Medical Research Institute – University of Queensland investigates molecular aberrations that could be used as biomarkers and/or therapeutic targets for late stage cancer.



Distinguished Professor Judith Clements studied Endocrinology at Monash University in Melbourne, Australia where she received her PhD in 1989. Since then she has been a Research Fellow, Senior Research Fellow and Principal Research Fellow of the Australian National Health and Medical Research Council. She is Head of the Cancer Program at the Institute of Health and Biomedical Innovation and Translational Research Institute at Queensland University of Technology and Scientific Director of the Australian Prostate Cancer Research Centre-Queensland. Her research interests are focused on mechanistic and translational studies of the role of the kallikrein-related serine peptidases in hormone dependent cancers, particularly prostate and ovarian cancers.